

# RELIANCE CALF MILK REPLACER



**RELIANCE CALF MILK REPLACER** is milk powder blend providing excellent calf response to the natural feed sources of milk fat, milk proteins and sugar.

## FEEDING RECOMMENDATIONS

**MIXING RATIO:** 125g **RELIANCE CALF MILK REPLACER** made up to one litre with water when fed twice daily, 250g Calf Milk Replacer made up to one litre with water when fed once daily.

Refer to the feeding recommendations and add required amount **of RELIANCE CALF MILK REPLACER** powder to half the volume of hot water (40-45°C), not boiling water. Mix briskly for 1 minute to a creamy smooth consistency and add cool water to make up the required amount (e.g. make up to 1L, do not add 1L water to 125g powder).

- Feed at body temperature (38-39°C).
- Mix only when required. Keep bag securely closed when not in use.
- Mixes easily and can be fed alone or added to milk to extend stocks or fortify.
- Feed after a good intake of colostrum has been achieved (from four days after birth).
- Ensure that clean drinking water is available at all times.
- Ensure that calves have access to hay and quality calf feed.
- Keep buckets, bottles and other equipment clean.

**STORAGE:** Store between 10 and 20°C in a clean, cool, dry and vermin-free environment. Do not expose to direct sunlight and avoid contact with floor and walls.

AGE IN DAYS	LITRES PER FEED	GRAMS PER DAY *	ONCE A DAY SYSTEM		TWICE A DAY SYSTEM	
			NO. FEEDS PER DAY	CONCENTRATION	NO. FEEDS PER DAY	CONCENTRATION
1 - 4	2.5	-	2	Colostrum Only	2	Colostrum Only
5 - 14	2	500	2	125 grams/litre	2	125 grams/litre
15 - 28	2.5	625	1	250 grams/litre	2	125 grams/litre
29 - weaning	2.5	625 -750	1	250 grams/litre	2	125 grams/litre

When higher rates of growth are preferred (accelerated growth programmes) or when calves have been stressed (e.g. disease recovery or environment), a higher rate of feeding may be applicable. Customised feeding plans are available by contacting your Reliance calf-rearing specialist.

## INGREDIENTS SELECTED FROM

Milk Powder, Vitamin A, D3, E, K3, B1, B2, B3, B5, B6, B12, Ascorbic Acid, Potassium, Lasalocid (Bovatec®), Niacin, Pantothenic Acid, Biotin, Cobalt (organic), Copper (organic), Dextrose, Ferrous Iron, Iodine, Selenium (organic), Magnesium, Manganese (organic), Zinc (organic), Antioxidants, Lecithin.

NOTE: MILK POWDER IS CASEIN BASED WHICH TYPICALLY CURDS BUT CURD STRENGTH MAY BE VARIABLE.

You can rely on Reliance

**RELIANCE**  
FEEDS

### TYPICAL ANALYSIS (APPROXIMATE ON AN AS FED BASIS)

Protein	25%
Fat	20%
Lactose	44%
Ash	6%

### COCCIDIOSTAT

#### **WARNING STATEMENTS**

**READ LABEL BEFORE USE**

**FOR ANIMAL TREATMENT ONLY**

**KEEP OUT OF THE REACH OF CHILDREN**

**NOT FOR HUMAN CONSUMPTION**

### Reliance Calf Milk Replacer with 500g/tonne Bovatec® 20CC.

**CONTAINS** Bovatec® 20CC which is registered pursuant to the ACVM Act 1997, No. A009679. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**INDICATIONS** Contains 100mg/kg of lasalocid sodium as 500g/tonne Bovatec® 20CC as an aid in the control of coccidiosis caused by Eimeria species in cattle.

Feed 500grams of Reliance Calf Milk Replacer per 50kg calf to provide 1mg of lasalocid per kg of body weight. Ensure young animals are adapted to a medicated feed at least 2-3 weeks prior to weaning as coccidial challenge is most likely to occur in the first eight weeks after weaning.

**DIRECTIONS FOR USE** This product must not be used for growth promotion in ruminants intended for human consumption. By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

**DOSE RATE** For coccidiosis control, minimum dose rate of 5mg Bovatec®/kg live weight (1mg lasalocid sodium/kg live weight) is required per day. Up to twice this dose rate is indicated where severe coccidial challenge is expected or a wide range of individual live weights occur within a group.

**CAUTION** Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec®, as ingestion by these species may be fatal. Do not feed to deer, milking goats, sheep, laying poultry and other non-ruminant animals.

#### **CONTRAINDICATIONS**

- Care must be exercised when feeding concurrently with other antimicrobials.
- Do not feed with other ionophores, e.g. monensin capsules, liquid or premix.
- Do not exceed the recommended dose rates.
- Do not feed more than a total of 400mg lasalocid sodium/head/day to lactating cattle.
- Not to be used for single-dose treatment.
- Disposal: avoid contamination of any water supply with this feed or its empty bag.

**WITHHOLDING PERIOD** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

#### **Cattle**

**Meat:** Nil

**Milk:** Cattle Nil

**Not to be used for bobby calves**

#### **MANUFACTURED FOR AND DISTRIBUTED BY:**

**Farmlands Nutrition**

Private Bag 1968, Dunedin 9054, New Zealand.